

Instructions for Patients Having Pan Facial Fat Grafting Surgery

2 Weeks Before Surgery

- Avoid all aspirin containing products, all anti-inflammatory medications (ie Ibuprofen such as Advil or Motrin), all products containing 400 IU's of vitamin E and all herbal products. These products can increase bleeding both during and after surgery and may lead to increased postoperative bruising, swelling and recovery time.
- Stop or at least cut down on smoking. Smoking constricts small blood vessels causing delayed healing, increased scarring and increased recovery time.
- Begin taking a multivitamin tablet daily, which can speed healing and shorten recovery time. It is recommended to continue your multivitamin for one month following surgery.
- Arrange for transportation to and from surgery. You must have someone responsible pick you up and remain with you for a minimum of 24 hours. You cannot take a taxi home on your own.
- It is recommended to restrict activity for approximately 2 weeks following surgery. It is advisable to make arrangements for assistance.
- Avoid dental appointments 2 weeks prior to your surgery as well as 2 weeks post operatively.
- Arnica capsules or pellets can be taken orally starting 2 days before surgery and can continue for 5 days post op. This homeopathic medication can assist in decreasing bruising and swelling due to surgery.

24 Hours Before Surgery

- Avoid alcohol and smoking.
- Nothing to eat or drink after midnight the night before your surgery (no water, no gum). If surgery is booked after 12 noon, clear fluids may be allowed up to 6 hours before surgery (ie: black coffee/tea, clear juice, water). Our nurse will advise you on this when you are given your time for surgery.

Day of Surgery

- Wear loose, comfortable button-up/zipper clothing (ie: jog suit, slip on shoes).
- Leave valuables at home (remove contact lenses, jewelry or piercing of any kind).

- Take any regular prescribed medications (ie. blood pressure medications) with a small amount of water.

After Surgery

- Some pain and nausea can be expected. This can be relieved with anti-pain and anti-nausea medications. Take them as directed. If you are coming from out of town, please have these medications with you.
- Maintain a regular fluid intake during the day. Gatorade is a good option.
Restrict fluids after supper; increased fluid intake will increase swelling.
- Rest is important to minimize pain, nausea and swelling.
- Keeping your head elevated after surgery will help decrease swelling.
- Begin a light nutritional regime and increase food intake as tolerated.
- Take Tylenol after surgery to decrease pain. 24 hours after surgery you may begin taking Advil.

**Call (403) 228-1313 if you have any questions or concerns.
If leaving a message please leave a number.**

Post-operative Day 1

- Be aware of any unusual swelling or a sudden increase in pain. If either of these are noted, phone the office as soon as possible.
- Continue to maintain an elevated position.
- Use cold compresses for 72 hours to decrease swelling.
- Be aware of decreased skin sensation. Do not apply excessive heat or cold to the operative areas.
- Do not smoke and rest as much as possible.

- A nurse will call you by noon to assess how you are doing. At this time, you will be given further post-operative instructions as well as an appointment for your first post-operative visit.